

Association between maternal lifestyle during early pregnancy and onset phase of childhood overweight

Kohta SUZUKI, Daisuke ANDO,
Taichiro TANAKA, Miri SATO,
Naoki KONDO, Zentaro YAMAGATA

Department of Health Sciences, School of
Medicine, University of Yamanashi

Background

- We previously clarified the association of maternal lifestyle during early pregnancy, including smoking and breakfast consumption, with childhood obesity and overweight at 5 years of age and 9–10years of age.
- To prevent childhood obesity, we considered that it was important to specify the risk factor about each onset phase.


Objectives

- We aimed to clarify the association between maternal lifestyle during early pregnancy and being childhood overweight between birth and 5 years of age, and between 5 years of age and 9–10 years of age.

Participants and Study Design

- The study population comprised children born between April 1, 1991 and March 31, 1997, in Japan, and their mothers.
- These subjects are the participants of Project Koshu, a dynamic ongoing prospective cohort study of pregnant women and their children in a Japanese rural area that commenced in 1988.


Project Koshu: A dynamic prospective cohort study



To ascertain the lifestyle habits of the expectant mothers, we conducted a questionnaire-based survey.

We surveyed the lifestyle habits of the children and their mothers by using a questionnaire.


During the checkups, we also obtained the growth and physical data of the children.



Outcomes of this study

- We classified the cases of the childhood overweight at 9–10 years of age to two groups below.
- Group 1: Being overweight between birth and 5 years of age
- Group 2: Being overweight between 5 years of age and 9–10 years of age
- Childhood overweight at 5 years of age and 9–10 years of age were defined as international cutoff value.



(Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000;320:1240-1243.)



5 years of age

9–10 years of age



Group 1



Overweight

Overweight

Group 2



Normal weight

Overweight

Statistical analysis

- Dependent variables: Group 1 or Group 2
- Independent variables:
 - Smoking habits during early pregnancy
 - Breakfast habits just before pregnancy
- We used a multiple logistic regression analysis to adjust the variables and the confounding factors, namely maternal age at delivery and maternal BMI at non-pregnancy.

Results - Participants




Table 1: Adjusted odds ratio (OR) and 95% confidence interval (CI) for maternal lifestyle factors that affected childhood overweight (Group1)

Lifestyle	Crude		Adjusted ^a	
	OR	95% CI	OR	95% CI
Smoking during early pregnancy				
Current smoker	4.15	(1.93 – 8.93)	3.56	(1.41 – 8.97)
Ex-smoker and Non-smoker				
Sleep duration				
Less than 8 h/d	1.78	(1.02 – 3.11)	1.52	(0.80 – 2.88)
More than 8 h/d				
Breakfast consumption				
I occasionally skip	2.18	(1.21 – 3.90)	2.51	(1.23 – 5.12)
I do not skip				
Maternal age at delivery			1.06	(0.99 – 1.15)
Maternal body mass index at non-pregnancy			1.21	(1.12 – 1.32)

a: Adjusted by maternal age, maternal body mass index, smoking status, sleep duration, and breakfast

Table 2: Adjusted odds ratio (OR) and 95% confidence interval (CI) for maternal lifestyle factors that affected childhood overweight (Group 2)

Lifestyle	Crude		Adjusted ^a	
	OR	95% CI	OR	95% CI
Smoking during early pregnancy				
Current smoker	1.01	(0.35 – 2.90)	0.76	(0.22 – 2.67)
Ex-smoker and Non-smoker				
Sleep duration				
Less than 8 h/d	0.93	(0.58 – 1.50)	0.80	(0.47 – 1.34)
More than 8 h/d				
Breakfast consumption				
I occasionally skip	1.10	(0.58 – 2.07)	1.04	(0.50 – 2.19)
I do not skip				
Maternal age at delivery			1.03	(0.96 – 1.10)
Maternal body mass index at non-pregnancy			1.14	(1.05 – 1.23)

a: Adjusted by maternal age, maternal body mass index, smoking status, sleep duration, and

Maternal lifestyle during pregnancy

5 years



Overweight

9–10 years




Overweight

Birth






Early onset



Normal weight

Late onset



The concept of Fetal Programming (Barker hypothesis)

Maternal under nutrition



Birth weight ↓



Weight gain during
early childhood ↑



Childhood and adult
overweight and obesity ↑



Type 2 diabetes,
hypertension, and CHD ↑

=

?

Maternal smoking

Maternal skipping
breakfast

Our data may represent
this relationship.



Moreover...

- These results suggested that late onset of childhood overweight was associated with other risk factors that were not considered in this study, such as childhood lifestyle including diet and exercise.
- For the prevention of childhood overweight, it may be important to carry out the continuous prevention program of mother and child which starts from pre-pregnancy.

Conclusion

- These results suggested that there were some differences of risk factor about each onset phase of childhood overweight.
- Especially, we believe that maternal lifestyle during and just before pregnancy is an important contributor to the prevention of being childhood overweight of early infant.

Acknowledgement

- Koshu project was conducted with the cooperation of the Koshu City administration office.
- We thank the public health nurses working at Koshu City for collaboratively implementing this study.