

SPATIAL DISTRIBUTION OF MORTALITY AND TEMPERATURE ALERT DAYS IN SPAIN DURING THE HEAT-WAVES

Diana Gómez-Barroso, Víctor M. Flores, Fernando Simón.

Ciberesp.Centro Nacional de Epidemiología, Instituto de Salud
Carlos III

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

1. INTRODUCTION

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

Since 20th century, heat has been associated with excess of mortality



Summer 2003

mayor Heat
wave



Increased
mortality

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

In Spain, summer 2003:

- High temperatures
- Long periods
- June, July and August

Reached a Peak → August → Notable increased general mortality affecting population over age 74 years

Geographical and temporal variability

July 2006 less intense Heat wave took place

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

2.OBJECTIVE

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

To compare the spatial distribution of mortality and temperature alert days in Spain during August 2003 and July 2006 using Kriging method.

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

2.METHODS

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

Information Sources:

Daily Mortality Monitory System of CNE

- Computerized Civil registries
- INE (National Statistic Institute)

Temperatures

- AEMET (State Agency for Meteorology)

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

Estimation Mortality Alert Days

Provincial capitals and towns of over 120.000 inhabitants

Excluded small towns and villages



Reduce random variability of mortality

Enhance alert specificity

Islands no correlated with mainland Spain

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

Estimation Mortality Alert Days

- Restricted age over 74 years
- Estimated expected mortality using time series:
 - 1997-2002 August 2003
 - 1998-2004 July 2006
- Variability and trend was corrected
- The expected number of deaths for each calendar day was estimated using the mean for that day and two days on either side of it, for all years included in the series.
- Used an adaptation of the Algorithmic Cumulative Sum method to estimate alert days.
- The alert threshold was set at the 90 percentile of the distribution of consecutive days of observed mortality exceeding expected during the previous two years.

ciberesp

Estimation Temperature Alert Days

Temperature alert day: maximum and minimum temperatures exceeded the 95 percentile of the maximum and minimum daily means for period 1971-2000

Spatial analysis

- Quantified mortality and temperature alert days in each town
- Results applied to centroid of each town
- Estimation of Risk by Ordinary Kriging
 - “Spatial estimation Method based on the spatial correlation among the values observed at different distances and directions(semivariogram) which enables a value to be interpolated for points where no information is available”(Opromolla)
- The Standard Error was also calculated.

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

METHOD

Ordinary Kriging

$$Z_{OK}^*(u_0) = \sum_{\alpha=1}^{n(u_0)} \lambda_{\alpha}^{OK}(u_0) Z(u_{\alpha})$$

Anisotropy, Exponential model

4 method for estimations:

- Ordinary least squares (OLS)
- Weighted least squares (WLS)
- Restricted maximum likelihood (REML)
- Maximum likelihood (MLE)

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

4. RESULTS

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

Towns:

59 in 2003


41 in 2006



Meteorological Stations:

89 in 2003

47 in 2006



AUGUST 2003

Estimation risk mortality alert days

Exponential model fitted by REML

Angle 40° NE_SW direction

Estimation risk temperature alert days

Exponential model fitted by MLE


Angle 45° NE_SW direction

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

Mortality Alerts Days August 2003

Temperatures Alerts Days, August 2003



JULY 2006

Estimation risk mortality alert days

Exponential model fitted by REML

Angle 60° NE_SW direction

Estimation risk temperature alert days

Exponential model fitted by MLE


Angle 45° NE_SW direction

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

Mortality Alerts Days July 2006

Temperatures Alerts Days, July 2006




5. LIMITATIONS

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

- Number of towns and high distance between them, involve a smaller degree of confidence in some areas



Standard error maps

- Mortality alert days estimations do not take in account the total numbers of deaths associated with heat.
- Temperature alert days estimations do not take in account degrees over the threshold
- Possible associated bias; Threshold for determining a mortality alert and temperature excess is not define for the same number of municipalities.

ciberesp

6. CONCLUSIONS

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública

- Possible association between the spatial distribution of risk of mortality alerts and risk of temperature alerts during periods of excessive heat.
- The longer and more intense the period of heat, the stronger this association seems to be.
- First study attempting to correlate mortality and temperatures alert days
- Multivariate methods: Cokriging and Regression Kriging
- This methods included in surveillance services can influence decision-making in Public Health.
- This the direction we are currently working on

ciberesp



Thanks

ciberesp

Centro de Investigación Biomédica en red
Epidemiología y Salud Pública